

National and International Childhood Cancer Incidence and Time Trends

Charles Stiller

**Childhood Cancer Research Group
Department of Paediatrics
University of Oxford
UK**

Childhood Cancer 2012


Childhood Cancer Incidence in the UK, 1996-2005

Age-standardized rate (ASR)	142 per million
Cumulative risk in first 15 years of life	1 in 484
Sex ratio (M/F)	1.2

Source: National Registry of Childhood Tumours

Annual Incidence by Single Year of Age All Cancers, Great Britain, 1991-2000

Fig 3.1 All cancers




Britain 1991-2000

Source: National Registry of Childhood Tumours

Stiller C (ed.) *Childhood Cancer in Britain: Incidence, Survival, Mortality*. OUP, 2007

Contributions of the 12 Main Groups of ICC-3* to Age-standardized Incidence, Great Britain, 1991-2000

Fig3.2 Age-standardised rate



Britain 1991-2000


***International Incidence of Childhood Cancer, Third Edition**
(Steliarova-Foucher E et al. *Cancer* 2005, **103**, 1457-1467)

Source: National Registry of Childhood Tumours

Stiller C (ed.) *Childhood Cancer in Britain: Incidence, Survival, Mortality*. OUP, 2007

Annual Incidence by Single Year of Age Lymphomas (ICCC-3 Group II), Great Britain, 1991-2000

Fig 3.14 Lymphomas and reticuloendothelial neoplasms




Britain 1991-2000

Source: National Registry of Childhood Tumours

Stiller C (ed.) *Childhood Cancer in Britain: Incidence, Survival, Mortality*. OUP, 2007

Annual Incidence by Single Year of Age Principal Embryonal Tumours, Great Britain, 1991-2000

Fig 3.21 Embryonal tumours




Britain 1991-2000

Source: National Registry of Childhood Tumours

Stiller C (ed.) *Childhood Cancer in Britain: Incidence, Survival, Mortality*. OUP, 2007

Childhood Cancer Incidence in Europe, 1988-1997



British Isles: Ireland, UK

East: Belarus, Estonia, Hungary, Slovakia

North: Denmark, Finland, Iceland, Norway


South: Italy, Malta, Slovenia, Spain, Turkey

West: France, Germany, Netherlands, Switzerland

Source: Automated Childhood Cancer Information System (ACCIS)

Stiller CA et al. *Eur J Cancer* 2006, **42**, 1952-1960.

Childhood Cancer Incidence in the Americas in the 1980s



Colombia: Cali


Costa Rica: national

Uruguay: national

USA: SEER registries

Parkin DM et al. (eds.) *International Incidence of Childhood Cancer, Volume 2*. IARC, 1998.


Childhood Cancer Incidence in the USA, 1999-2003



Source: 38 SEER & NPCR registries

Howe HL et al. *Cancer* 2006, **107**, 1711-1742.

Childhood Cancer Incidence in Asia & Oceania in the 1980s



Hong Kong, Israel, Australia: National


India: Bombay

Japan: 5 registries

Thailand: 4 registries

Parkin DM et al. (eds.) *International Incidence of Childhood Cancer, Volume 2*. IARC, 1998.

Childhood Cancer Incidence in Africa in the 1990s



Algeria: 4 registries


Tunisia: 3 registries

Nigeria: Ibadan

Uganda: Kyadondo County

Zimbabwe: Harare (Africans)

Ewing Sarcoma of Bone: Incidence in the 1980s



Australia, Germany, Hong Kong: National

UK: England & Wales

Japan: 5 registries

USA: SEER registries

Parkin DM et al. (eds.) *International Incidence of Childhood Cancer, Volume 2*. IARC, 1998.

Adrenocortical Carcinoma in Southern Brazil


	ASR per million	
	Incidence	Mortality
São Paulo (1969-1978) ¹	1.5	
Goiânia (1989-1994) ²	2.8	
Curitiba (1998-2003) ³	3.5?	1.6
Other world regions ²	<0.5	

1 Parkin DM et al. (eds.) *International Incidence of Childhood Cancer*. IARC, 1988.

2 Parkin DM et al. (eds.) *International Incidence of Childhood Cancer, Volume 2*. IARC, 1998.

3 Pianovski MAD et al. *Pediatr Blood Cancer* 2006,**47**,56-60

Malignant Melanoma: Incidence in the 1980s




Australia, New Zealand: National

UK: England & Wales

USA: SEER registries

All Childhood Cancer, Great Britain, 1966-2005

Age-sex-standardized rates by 5-year period of diagnosis




Annual average % change (95% CI) 1.0 (0.9,1.1)

Source: National Registry of Childhood Tumours

All Childhood Cancer, Great Britain, 1966-2005


Age-sex-standardized rates by 5-year period of diagnosis



	Annual average % change (95% CI)
Leukaemia (ICCC-3 I)	0.7 (0.6,0.9)
CNS tumours (ICCC-3 III, Xa)	1.3 (1.2,1.5)
All other childhood cancers	1.0 (0.9,1.1)


Source: National Registry of Childhood Tumours

Childhood Leukaemia in England and Wales, 1911-1960



Mortality at age 0, 1-4, 5-9, 10-14

Acute Lymphoblastic Leukaemia in Europe, 1970-1999




West: Denmark, Finland, France, Germany, Iceland, Ireland, Italy, Malta, Netherlands, Norway, Spain, Switzerland, UK

East: Belarus, Estonia, Slovakia, Slovenia, Turkey

Source: ACCIS

Steliarova-Foucher E et al. *Lancet* 2004, **364**, 2097-2105

Primary Malignant Brain Tumours in the USA, 1973-1994




Age-standardized annual rates:	Δ Incidence		X Mortality		
	1973	1977	1982	1984	1985
No. of CT scanners	8	921			
No. of MRI units			10	108	371

Source: SEER Program


Smith MA et al. *J Natl Cancer Inst* 1998, **90**, 1269-1277

Childhood CNS Tumours in Great Britain, 1966-2005




Source: National Registry of Childhood Tumours

Childhood Cancer Incidence in Kyadondo County, Uganda 1960-1971 and 1991-1997




Hepatocellular Carcinoma in Taiwan, 1981-1994

Incidence & mortality (age 6-14)



Hepatitis B mass vaccination began 1984

Thyroid Cancer Incidence in Children and Adolescents Ukraine, 1989-2008




ASR at age 0-19 in 1989, 1990-1994, 1995-1999, 2000-2004, 2005-2008

Regions classified by reconstructed ^{131}I thyroid dose

High exposure: >35 mGy



Low exposure: ≤ 35 mGy

Cutaneous Malignant Melanoma in Children & Adolescents Sweden, 1973-2002



ASR at age 0-19

Childhood Melanoma in Australia, 1983-2006



Observed incidence by year of diagnosis

Underlying trends


1983-1993 +8.4% per year

1993-2006 -8.5% per year


Neuroblastoma in Japan

Incidence, mortality and survival, 1970-1994

Osaka Prefecture excluding Osaka City



Neuroblastoma in Europe, 1988-1997



Incidence at age 0, 1-4, 5-9, 10-14

British Isles: Ireland, UK

East: Belarus, Estonia, Hungary, Slovakia

North: Denmark, Finland, Iceland, Norway

South: Italy, Malta, Slovenia, Spain, Turkey

West: France, Germany, Netherlands, Switzerland

Source: ACCIS

Spix C et al. *Eur J Cancer* 2006, **42**, 2081-2091.

Future Work

International Incidence of Childhood Cancer, Volume 3 (IICC-3)

ACCIS-2

- **Calls for data have been issued**
- **Years of diagnosis to at least 2005**
- **Age 0-19 in both studies**
- **Increased geographical coverage, especially outside
W Europe, N America and Australia / New Zealand**

Future Trends

Most marked trends will probably occur in lower income countries

- **Kaposi sarcoma in areas of high HHV8 expected to rise and fall with childhood HIV**
- **Hepatocellular carcinoma in areas of moderately high incidence should decline with uptake of hepatitis B vaccination**
- **Burkitt lymphoma may increase if geographical range of holoendemic malaria widens with climate change**
- **Recorded incidence of many other cancers may rise as populations become more affluent and availability of facilities for diagnosis and treatment increases**